



CHALLENGING SFA/POP CASE 1

- 77 Y-OLD
- EX HEAVY SMOKER
- PREVIOUS FEM-POP VEIN BYPASS
BECAUSE OF 2nd TOE necrosis (2019)
- TRAUMA 1ST TOE WITH INFECTION
AND NECROSIS



RT

RT

RT



RAO 37

RAO 27

APPROACH

Three Radiopaque Markers

- Markers facilitate accurate assessment of position
- Unique spacing provides easy measurement of common stent and balloon sizes
- Embedded marker is 1 mm from distal tip; 40 mm and 60 mm spacing


Minimum Sheath Compatibility: 4 Fr

- Intraluminal 0.035
- Intraluminal 0.018
- Subintimal 0.035
- Open surgery

RT







RT





RAO 35 deg


ASAHI GLADIUS MG PV ES


















[> J Endovasc Ther. 2018 Jun;25\(3\):334-342. doi: 10.1177/1526602818763352. Epub 2018 Mar 20.](#)

Endovascular Treatment of Severely Calcified Femoropopliteal Lesions Using the "Pave-and-Crack" Technique: Technical Description and 12-Month Results

[Marina Dias-Neto ^{1 2}, Manuela Matschuck ³, Yvonne Bausback ³, Ursula Banning-Eichenseher ³, Sabine Steiner ³, Daniela Branzan ⁴, Holger Staab ⁴, Ramon L Varcoe ^{5 6}, Dierk Scheinert ³, Andrej Schmidt ³](#)

